

The Cardy-Verlinde Formula and Asymptotically Flat Charged Black Holes

Donam Youm¹

ICTP, Strada Costiera 11, 34014 Trieste, Italy

Abstract

We show that the modified Cardy-Verlinde formula without the Casimir effect term is satisfied by asymptotically flat charged black holes in arbitrary dimensions. Thermodynamic quantities of the charged black holes are shown to satisfy the energy-temperature relation of a two-dimensional CFT, which supports the claim in our previous work (Phys. Rev. D61, 044013, hep-th/9910244) that thermodynamics of charged black holes in higher dimensions can be effectively described by two-dimensional theories. We also check the Cardy formula for the two-dimensional black hole compactified from a dilatonic charged black hole in higher dimensions.

May, 2001

¹E-mail: youmd@ictp.trieste.it

Recently, thermodynamics of the holographic duals of various black hole solutions have been actively studied, after Verlinde proposed [1, 2] that the Cardy formula [3] for the two-dimensional conformal field theory (CFT) can be generalized to arbitrary spacetime dimensions $D - 1$ in the form

$$S = \frac{2\pi R}{D-2} \sqrt{E_c(2E - E_c)}, \quad (1)$$

the so-called Cardy-Verlinde formula [1, 2], where R is the radius of the system, E is the total (thermal excitation) energy and E_c , called the Casimir energy, is the subextensive part of E . (Cf. The quantum effects to the Cardy-Verlinde formula were studied in Refs. [4, 5].) So far, mostly asymptotically AdS black hole solutions have been considered [6, 7, 8, 9, 10]. In Ref. [10], it is shown that even the Schwarzschild and the Kerr black hole solutions, which are asymptotically flat, satisfy the modification of the Cardy-Verlinde formula,

$$S = \frac{2\pi R}{D-2} \sqrt{2E_c E}, \quad (2)$$

to the case of the ground state with zero conformal weight. In this note, we show that this result holds also for various charged black hole solutions with asymptotically flat spacetime.

First, we consider the single-charged dilatonic black hole solution with an arbitrary dilaton coupling parameter a in arbitrary spacetime dimensions D . The action is

$$S_D = \frac{1}{2\kappa_D^2} \int d^D x \sqrt{-G} \left[\mathcal{R} - \frac{4}{D-2} (\partial\phi)^2 - \frac{1}{4} e^{2a\phi} F^2 \right], \quad (3)$$

where κ_D is the D -dimensional gravitational constant, ϕ is the dilaton field and F is the field strength of the $U(1)$ gauge potential $A = A_M dx^M$ ($M = 0, 1, \dots, D-1$). The nonextreme black hole solution to the field equations of this action is given by

$$\begin{aligned} G_{MN} dx^M dx^N &= -H^{-\frac{4(D-3)}{(D-2)\Delta}} f dt^2 + H^{\frac{4}{(D-2)\Delta}} \left[f^{-1} dr^2 + r^2 d\Omega_{D-2}^2 \right], \\ e^\phi &= H^{\frac{(D-2)a}{2\Delta}}, \quad A_t = \frac{2}{\sqrt{\Delta}} \frac{m \sinh \alpha \cosh \alpha}{r^{D-3}} H^{-1}, \end{aligned} \quad (4)$$

where

$$\begin{aligned} H &= 1 + \frac{m \sinh^2 \alpha}{r^{D-3}}, \quad f = 1 - \frac{m}{r^{D-3}}, \\ \Delta &= \frac{(D-2)a^2}{2} + \frac{2(D-3)}{D-2}. \end{aligned} \quad (5)$$

For special values of Δ , this solution can be realized in string theories, e.g., the $\Delta = 4/n$ case with a positive integer n can be obtained by compactifying intersecting n numbers of branes with equal charges. The $a = 0$ case is the Reissner-Nordström black hole.

The ADM mass M , the Bekenstein-Hawking entropy S , the Hawking temperature T , the electric charge Q , and the chemical potential Φ of the solution are

$$\begin{aligned} M &= \frac{V_{D-2}}{2\kappa_D^2} \left[\frac{4(D-3)}{\Delta} m \sinh^2 \alpha + (D-2)m \right], \\ S &= \frac{4\pi V_{D-2}}{2\kappa_D^2} m^{\frac{D-2}{D-3}} \cosh^{\frac{4}{D-3}} \alpha, \quad T = \frac{D-3}{4\pi} m^{-\frac{1}{D-3}} \cosh^{-\frac{4}{D-3}} \alpha, \\ Q &= \frac{V_{D-2}}{2\kappa_D^2} \frac{2(D-3)}{\sqrt{\Delta}} m \sinh \alpha \cosh \alpha, \quad \Phi = \frac{2}{\sqrt{\Delta}} \tanh \alpha, \end{aligned} \quad (6)$$

where $V_n = 2\pi^{\frac{n+1}{2}}/\Gamma\left(\frac{n+1}{2}\right)$ denotes the volume of the unit S^n . These thermodynamic quantities satisfy the first law of black hole thermodynamics:

$$dM = TdS + \Phi dQ. \quad (7)$$

According to the holographic principle, the thermodynamic quantities of a black hole solution can be identified with the corresponding thermodynamic quantities of the holographic dual theory [11]. In the case of a charged black hole solution, the total energy $E_{tot} = M$ of the dual theory can be separated into two parts: the contribution of the supersymmetric background (i.e., the zero temperature energy) and that of thermal excitation. Following Refs. [6, 10], we define the zero temperature contribution, called the proper internal energy, as

$$E_q = \frac{V_{D-2}}{2\kappa_D^2} \frac{4(D-3)}{\Delta} m \sinh^2 \alpha. \quad (8)$$

So, the thermal excitation energy is

$$E = E_{tot} - E_q = \frac{V_{D-2}}{2\kappa_D^2} (D-2)m. \quad (9)$$

If we assume that the holographic dual theory is conformal, then the pressure $p = -\left(\frac{\partial E}{\partial V}\right)_{S,Q}$ is given by

$$p = \frac{E}{(D-2)V} = \frac{V_{D-2}}{2\kappa_D^2} \frac{m}{V}, \quad (10)$$

where V is the volume of the system. So, the Casimir energy of the holographic dual theory is given by

$$E_c = (D-2)(E_{tot} + pV - TS - \Phi Q) = \frac{V_{D-2}}{2\kappa_D^2} 2(D-2)m. \quad (11)$$

Since $E_c = 2E$, the Cardy-Verlinde formula (1) would yield $S = 0$. So, we have to consider the modified Cardy-Verlinde formula (2), valid for the case of the ground state with zero conformal weight. It is straightforward to show that thermodynamic

quantities S , E and E_c , respectively given in Eqs. (6,9,11), satisfy Eq. (2), provided we identify the length scale of the system as

$$R = m^{\frac{1}{D-3}} \cosh^{\frac{4}{D}} \alpha. \quad (12)$$

Just like the Schwarzschild black hole case [10], total thermal excitation energy (9) can be put into the following suggestive form:

$$E = \frac{c}{6} \pi^2 R \left(\frac{2T}{D-3} \right)^2, \quad (13)$$

with the effective central charge c given by

$$\frac{c}{6} = 2E_c R = \frac{V_{D-2}}{2\kappa_D^2} 4(D-2) m^{\frac{D-2}{D-3}} \cosh^{\frac{4}{D}} \alpha = \frac{D-2}{\pi} S. \quad (14)$$

This is the energy-temperature relation of a two-dimensional CFT with the characteristic length $R = m^{\frac{1}{D-3}} \cosh^{\frac{4}{D}} \alpha$ and effective temperature $\tilde{T} = 2T/(D-3)$ with T given in Eq. (6). Note, $c \propto S$, which is in accordance with the holographic principle. We can write Eq. (13) also in the form

$$ER = \frac{c}{24}, \quad (15)$$

which is the ground state energy of a two-dimensional CFT. The fact that thermodynamic quantities of a charged black hole solution (4) satisfy the relation (13) for a two-dimensional CFT supports the claim in our previous work [12] that thermodynamics of higher-dimensional charged black hole solutions can be effectively described by two-dimensional theories.

It is straightforward to show that the modified Cardy-Verlinde formula (2) is satisfied also by multi-charged black hole solutions in string theory. Thermodynamic quantities of such solutions are generally given by

$$\begin{aligned} M &= \frac{V_{D-2}}{2\kappa_D^2} \left[(D-3)m \sum_i \sinh^2 \alpha_i + (D-2)m \right], \\ S &= \frac{4\pi V_{D-2}}{2\kappa_D^2} m^{\frac{D-2}{D-3}} \prod_i \cosh \alpha_i, \quad T = \frac{D-3}{4\pi} m^{-\frac{1}{D-3}} \prod_i \cosh^{-1} \alpha_i, \\ Q_i &= \frac{V_{D-2}}{2\kappa_D^2} m \sinh \alpha_i \cosh \alpha_i, \quad \Phi_i = \tanh \alpha_i, \end{aligned} \quad (16)$$

where $i = 1, \dots, 4$ for $D = 4$ and $i = 1, 2$ for $D \geq 5$. These thermodynamic quantities satisfy the first law of black hole thermodynamics:

$$dM = TdS + \Phi_i dQ_i. \quad (17)$$

We define the proper internal energy as

$$E_q = \frac{V_{D-2}}{2\kappa_D^2} (D-3)m \sum_i \sinh^2 \alpha_i. \quad (18)$$

Then, the thermal excitation energy $E = E_{tot} - E_q$ is still given by Eq. (9). By using the assumed relation $pV = E/(D-2)$, which holds for the conformal holographic dual theory, we obtain the following Casimir energy:

$$E_c = (D-2)(E_{tot} + pV - TS - \Phi_i Q_i) = \frac{V_{D-2}}{2\kappa_D^2} 2(D-2)m. \quad (19)$$

So, the modified Cardy-Verlinde formula (2) is satisfied by the above thermodynamic quantities, provided we identify the length scale of the system as

$$R = m^{\frac{1}{D-3}} \prod_i \cosh \alpha_i. \quad (20)$$

It is straightforward to show that the thermal excitation energy E can also be put into the suggestive form (13) with R and T respectively given in Eqs. (20,16) and the effective central charge c given by

$$\frac{c}{6} = 2E_c R = \frac{V_{D-2}}{2\kappa_D^2} 2(D-2)m^{\frac{D-2}{D-3}} \prod_i \cosh \alpha_i = \frac{D-2}{\pi} S. \quad (21)$$

In our previous work [13], we showed that by compactifying the dual-frame action for the Hodge-dual action of Eq. (3) on a $(D-2)$ -sphere with the radius $\hat{\mu} = (m \sinh^2 \alpha)^{1/(D-3)}$ we obtain the following two-dimensional effective action:

$$S_2 = \frac{1}{2\kappa_2^2} \int d^2x \sqrt{-g} e^{\delta\phi} [\mathcal{R}_g + \gamma(\partial\phi)^2 + \Lambda], \quad (22)$$

where

$$\delta \equiv -\frac{D-2}{D-3}a, \quad \gamma \equiv \frac{D-1}{D-2}\delta^2 - \frac{4}{D-2}, \quad \Lambda = \frac{D-3}{2\hat{\mu}^2} \left[2(D-2) - \frac{4(D-3)}{\Delta} \right]. \quad (23)$$

The Weyl rescaling $\bar{g}_{\mu\nu} = \Phi^{\frac{\gamma}{\delta^2}} g_{\mu\nu}$ ($\Phi \equiv e^{\delta\phi}$) brings this action to the form:

$$S_2 = \frac{1}{2\kappa_2^2} \int d^2x \sqrt{-\bar{g}} [\Phi \mathcal{R}_{\bar{g}} + \Phi^{1-\frac{\gamma}{\delta^2}} \Lambda]. \quad (24)$$

The general static solution to the equations of motion for this action in the Schwarzschild gauge is

$$\begin{aligned} d\bar{s}^2 &= - \left[\frac{\delta^2}{2\delta^2 - \gamma} \left(\frac{x}{\ell} \right)^{2-\frac{\gamma}{\delta^2}} - 2\ell\kappa_2^2 \bar{M} \right] d\tau^2 + \left[\frac{\delta^2}{2\delta^2 - \gamma} \left(\frac{x}{\ell} \right)^{2-\frac{\gamma}{\delta^2}} - 2\ell\kappa_2^2 \bar{M} \right]^{-1} dx^2, \\ \Phi &= \frac{x}{\ell}, \end{aligned} \quad (25)$$

where $\ell \equiv 1/\sqrt{\Lambda}$ and $\bar{M}$ is the ADM mass of the solution. Thermodynamic quantities of the solutions are

$$\bar{M} = \frac{\delta^2}{2\delta^2 - \gamma} \frac{1}{2\ell\kappa_2^2} \Phi_H^{2-\frac{\gamma}{\delta^2}}, \quad \bar{T} = \frac{1}{4\pi\ell} \Phi_H^{1-\frac{\gamma}{\delta^2}}, \quad \bar{S} = \frac{2\pi}{\kappa_2^2} \Phi_H, \quad (26)$$

where Φ_H is the value of Φ at the event horizon, i.e., $\bar{g}_{\tau\tau}(\Phi_H) = 0$. If we assume the two dimensional system under consideration to be conformal, then $\bar{p}\bar{V} = \bar{M}$, so the Casimir energy is given by

$$\bar{M}_c = \bar{M} + \bar{p}\bar{V} - \bar{T}\bar{S} = \frac{\gamma}{2\delta^2 - \gamma} \frac{1}{2\ell\kappa_2^2} \Phi_H^{2-\frac{\gamma}{\delta^2}}. \quad (27)$$

These thermodynamic quantities satisfy the following Cardy formula [3] for a two-dimensional CFT:

$$\bar{S} = 2\pi \sqrt{\frac{c}{6} \left(L_0 - \frac{c}{24} \right)}, \quad (28)$$

with $L_0 = 2\ell\bar{M}$ and $c/6 = 4\ell\bar{M}_c$, provided $\gamma = \delta^2$. This is in agreement with the result in our previous work [13] that the action (24) is conformal if $\gamma = \delta^2$.

References

- [1] E. Verlinde, “On the holographic principle in a radiation dominated universe,” hep-th/0008140.
- [2] I. Savonije and E. Verlinde, “CFT and entropy on the brane,” hep-th/0102042.
- [3] J.L. Cardy, “Operator content of two-dimensional conformally invariant theories,” Nucl. Phys. **B270** (1986) 186.
- [4] S. Nojiri and S.D. Odintsov, “AdS/CFT correspondence, conformal anomaly and quantum corrected entropy bounds,” hep-th/0011115.
- [5] S. Nojiri and S.D. Odintsov, “AdS/CFT and quantum-corrected brane entropy,” hep-th/0103078.
- [6] R. Cai, “The Cardy-Verlinde formula and AdS black holes,” hep-th/0102113.
- [7] A.K. Biswas and S. Mukherji, “Holography and stiff-matter on the brane,” JHEP **0103** (2001) 046, hep-th/0102138.
- [8] D. Birmingham and S. Mokhtari, “The Cardy-Verlinde formula and Taub-Bolt-AdS spacetimes,” hep-th/0103108.
- [9] D. Klemm, A.C. Petkou and G. Siopsis, “Entropy bounds, monotonicity properties and scaling in CFTs,” hep-th/0101076.
- [10] D. Klemm, A.C. Petkou, G. Siopsis and D. Zanon, “Universality and a generalized C-function in CFTs with AdS duals,” hep-th/0104141.
- [11] E. Witten, “Anti-de Sitter space, thermal phase transition, and confinement in gauge theories,” Adv. Theor. Math. Phys. **2** (1998) 505, hep-th/9803131.
- [12] D. Youm, “Black hole thermodynamics and two-dimensional dilaton gravity theory,” Phys. Rev. **D61** (2000) 044013, hep-th/9910244.
- [13] D. Youm, “(Generalized) conformal quantum mechanics of 0-branes and two-dimensional dilaton gravity,” Nucl. Phys. **B573** (2000) 257, hep-th/9909180.